

S8 Table. Plasmids used in this study.

Plasmid name	Description	Reference
pJG1100	Suicide vector for mutant construction, Hyg ^R , Kan ^R , <i>sacB</i> .	[6]
pGA44	Integrative vector at L5 <i>attB</i> site, PTR promoter, Str ^R /Spect ^R .	[7]
pGA80	pMV261-derived vector, carrying the L5 <i>int</i> gene for expression <i>in trans</i> , lacking oriM, Kan ^R .	[7]
pJG1100- <i>espL</i> -UP/DOWN	Suicide vector for mutant construction derived from pJG1100, carrying 1 kb upstream and downstream regions of <i>espL</i> .	This study
pGA44- <i>espL</i>	Vector for complementation of $\Delta espL$ mutant strain. Derived from pGA44, <i>espL</i> is expressed by the PTR promoter.	This study
pGA44- <i>whiB6</i>	Vector for expression of <i>whiB6</i> . Derived from pGA44, <i>whiB6</i> is expressed by the PTR promoter.	This study
pGA44- <i>espL</i> .HA	Vector for complementation of $\Delta espL$ mutant strain. Derived from pGA44, <i>espL</i> .HA (tag at C-terminus) is expressed by the PTR promoter.	This study
pGA44-HA. <i>espL</i>	Vector for complementation of $\Delta espL$ mutant strain. Derived from pGA44, HA. <i>espL</i> (tag at N-terminus) is expressed by the PTR promoter.	This study
pGA44- <i>espE</i> .HA	Vector for expression of <i>espE</i> .HA (tag at C-terminus). Derived from pGA44, <i>espE</i> .HA is expressed by the PTR promoter.	This study
pGA44- <i>espE</i> .HA+ <i>pmycP1-espL</i>	Vector for expression of <i>espE</i> .HA (tag at C-terminus). Derived from pGA44, <i>espE</i> .HA is expressed by the PTR promoter. <i>espL</i> is expressed by the <i>mycP1</i> promoter.	This study

pGA44- <i>espE</i> .HA+ <i>pmycP1-whiB6</i>	Vector for expression of <i>espE</i> .HA (tag at C-terminus). Derived from pGA44, <i>espE</i> .HA is expressed by the PTR promoter. <i>whiB6</i> is expressed by the <i>mycP1</i> promoter.	This study
---	---	------------

Bibliography

1. Cortes T, Schubert OT, Rose G, Arnvig KB, Comas I, Aebbersold R, et al. Genome-wide Mapping of Transcriptional Start Sites Defines an Extensive Leaderless Transcriptome in *Mycobacterium tuberculosis*. *Cell Rep*. 2013;5: 1121–1131. doi:10.1016/j.celrep.2013.10.031
2. Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, et al. Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature*. 1998;393: 537–544. doi:10.1038/31159
3. Bottai D, Majlessi L, Simeone R, Frigui W, Laurent C, Lenormand P, et al. ESAT-6 secretion-independent impact of ESX-1 genes *espF* and *espG1* on virulence of *Mycobacterium tuberculosis*. *J Infect Dis*. 2011;203: 1155–1164. doi:10.1093/infdis/jiq089
4. Chen JM, Boy-Röttger S, Dhar N, Sweeney N, Buxton RS, Pojer F, et al. *EspD* is critical for the virulence-mediating ESX-1 secretion system in *Mycobacterium tuberculosis*. *J Bacteriol*. 2012;194: 884–893. doi:10.1128/JB.06417-11
5. Lou Y, Rybniker J, Sala C, Cole ST. *EspC* forms a filamentous structure in the cell envelope of *Mycobacterium tuberculosis* and impacts ESX-1 secretion: Filamentous structure formation by *EspC*. *Mol Microbiol*. 2017;103: 26–38. doi:10.1111/mmi.13575
6. Gomez JE, Bishai WR. *whmD* is an essential mycobacterial gene required for proper septation and cell division. *Proc Natl Acad Sci U S A*. 2000;97: 8554–8559. doi:10.1073/pnas.140225297
7. Kolly GS, Boldrin F, Sala C, Dhar N, Hartkoorn RC, Ventura M, et al. Assessing the essentiality of the decaprenyl-phospho- D -arabinofuranose pathway in *Mycobacterium tuberculosis* using conditional mutants: Druggability of the *M. tuberculosis* DPA pathway. *Mol Microbiol*. 2014;92: 194–211. doi:10.1111/mmi.12546